



# Edge irregular reflexive labeling on sun graph and corona of cycle and null graph with two vertices

Irfan Setiawan, Diari Indriati

*Department of Mathematics, Faculty of Mathematics and Natural Sciences, Universitas Sebelas Maret*

sunirfan74@gmail.com, diari\_indri@yahoo.co.id

---

## Abstract

Let  $G(V, E)$  be a simple and connected graph which set of vertices is  $V$  and set of edges is  $E$ . Irregular reflexive  $k$ -labeling  $f$  on  $G(V, E)$  is assignment that carries the numbers of integer to elements of graph, such that the positive integer  $\{1, 2, 3, \dots, k_e\}$  assignment to edges of graph and the even positive integer  $\{0, 2, 4, \dots, 2k_v\}$  assignment to vertices of graph. Then, we called as edge irregular reflexive  $k$ -labelling if every edges has different weight with  $k = \max\{k_e, 2k_v\}$ . Besides that, there is definition of reflexive edge strength of  $G(V, E)$  denoted as  $res(G)$ , that is a minimum  $k$  that using for labeling  $f$  on  $G(V, E)$ . This paper will discuss about edge irregular reflexive  $k$ -labeling for sun graph and corona of cycle and null graph, denoted by  $C_n \odot N_2$  and make sure about their reflexive edge strengths.

*Keywords:* Edge irregular reflexive labeling, reflexive edge strength, sun graph, corona of cycle and null graph

Mathematics Subject Classification: 05C78

DOI: 10.19184/ijc.2021.5.1.5

---

Received: 24 August 2020, Revised: 24 January 2021, Accepted: 29 April 2021.

## 1. Introduction

All graph discuss on this paper are simple, connected, and undirected. First, let simplified  $G(V, E)$  become  $G$ , vertex of graph denoted as  $v$ , and edge of graph denoted as  $e$  for easily on investigation. By definition from Wallis and Marr [10], graph labeling is a map that carries graph elements to numbers, usually to positive integer. The kinds of graph labeling divided as vertex-labelings, edge-labelings, and total labeling (vertex and edge-labelings). Based on survey that launch by Galian [5], there are many kinds of graph labelling. One of them is irregular total  $k$ -labeling.

According to Bača *et al.*[8], irregular total  $k$ -labeling divided be two kinds, that is edge irregular total  $k$ -labeling and vertex irregular total  $k$ -labeling. In 2017, Ryan *et al.* [6] has introduced new concept about irregular total  $k$ -labeling, that is vertex irregular reflexive total  $k$ -labeling and edge irregular reflexive total  $k$ -labeling. We call edge irregular reflexive total  $k$ -labeling if labeling  $f$  on  $G$  carries the positive integers 1 until  $k_e$  to edges of graph and carries the even positive integers 2 until  $2k_v$  to vertices of graph with  $k = \max\{k_e, 2k_v\}$ . The other specifications edge irregular reflexive total  $k$ -labeling is all of edges have different weight. The mapping positive integers to edge  $xy$  of graph denoted as  $f(xy)$  and the mapping even positive integers to vertex  $x$  of graph denoted as  $f(x)$ . Moreover, the weight edge  $xy$  of graph denoted as  $wt(xy)$ , where  $wt(xy) = f(x) + f(xy) + f(y)$ .

This paper also make investigate about reflexive edge strength of  $G$ , denoted as  $res(G)$ . To determine a lower bound of  $res(G)$  Ryan [6] gave a *Lemma 1.1* for all graph  $G$ ,

### Lemma 1.1.

$$res(G) \geq \begin{cases} \lceil \frac{|E(G)|}{3} \rceil, & \text{if } |E(G)| \not\equiv 2, 3 \pmod{6}, \\ \lceil \frac{|E(G)|}{3} \rceil + 1, & \text{if } |E(G)| \equiv 2, 3 \pmod{6}. \end{cases}$$

Some of  $res(G)$  has been determined, such as prisms graph  $D_n$  [2], cycle  $C_n$  [7], wheels graph  $W_n$  [2], corona of path and other graph,  $P_n \odot K_1$  and  $P_n \odot P_2$  [1], and other. This paper will investigate about sun graph and corona of cycle and null graph  $N_2$ .

## 2. The sun graph

Based on definition of sun graph by Wallis and Marr [10], an  $n$ -sun is a cycle  $C_n$  with an edge terminating in a vertex of degree 1 attached to each vertex and by Boulet [9] denoted as  $Sun_n$ . Then we will denote vertices in the cycle as  $x_i$  and pendant vertices denoted as  $y_i$ . So set of vertices  $V(sun_n) = \{x_i, y_i : 1 \leq i \leq n\}$ , consequently set of edges  $E(sun_n) = \{x_i x_{i+1}, x_i y_i : 1 \leq i \leq n\}$ . As a result  $sun_n$  has  $2n$  edges and vertices.  $Res(G)$  of  $sun_n$  can be obtained by *Theorem 2.1*.

**Theorem 2.1.** For  $sun_n$  with  $n \geq 3$ ,

$$res(sun_n) \doteq \begin{cases} 3, & \text{if } n \doteq 3, \\ 2\lceil \frac{n}{3} \rceil, & \text{if } n > 3. \end{cases}$$

*Proof.* First, we prove the lower bound of  $res(sun_n)$ . Since  $|E|$  of  $sun_n$  is  $2n$ , then by Lemma 1.1 we get

$$res(sun_n) \geq \begin{cases} \lceil \frac{2n}{3} \rceil, & \text{if } 2n \not\equiv 2, 3 \pmod{6}, \\ \lceil \frac{2n}{3} \rceil + 1, & \text{if } 2n \equiv 2, 3 \pmod{6}. \end{cases} \quad (1)$$

Let's prove the condition if  $n \doteq 3$ ,

Graph  $sun_3$  has 6 vertices and 6 edges. By (1) we get the lower bound of  $res(sun_3) = 2$ . Let's assume  $res(sun_3) = 2$ , we get maximum label of vertex and label of edges is 2, consequently the possibility of vertices and edges label for 6 edges  $x_i y_i (1 \leq i \leq 6)$  with the edge weight from 1 until 6 are,

$$\begin{aligned} wt_f(x_1 y_1) &= f(x_1) + f(x_1 y_1) + f(y_1) = 0 + 1 + 0 = 1 \\ wt_f(x_2 y_2) &= f(x_2) + f(x_2 y_2) + f(y_2) = 0 + 2 + 0 = 2 \\ wt_f(x_3 y_3) &= f(x_3) + f(x_3 y_3) + f(y_3) = 0 + 1 + 2 = 3 \\ wt_f(x_4 y_4) &= f(x_4) + f(x_4 y_4) + f(y_4) = 0 + 2 + 2 = 4 \\ wt_f(x_5 y_5) &= f(x_5) + f(x_5 y_5) + f(y_5) = 2 + 1 + 2 = 5 \\ wt_f(x_6 y_6) &= f(x_6) + f(x_6 y_6) + f(y_6) = 2 + 2 + 2 = 6 \end{aligned}$$

But, it's form can't be applied. Then, let's make assume that  $res(sun_n) = 3$ , we get  $\max f(x_i) = 2$  and  $\max f(x_i y_i) = 3$  for  $1 \leq i \leq 6$ , consequently the possibility label of vertices and edges as follows,

$$\begin{aligned} wt_f(x_1 y_1) &= f(x_1) + f(x_1 y_1) + f(y_1) = 0 + 1 + 0 = 1 \\ wt_f(x_2 y_2) &= f(x_2) + f(x_2 y_2) + f(y_2) = 0 + 2 + 0 = 2 \\ wt_f(x_3 y_3) &= f(x_3) + f(x_3 y_3) + f(y_3) = 0 + 1 + 2 = 3 \\ wt_f(x_4 y_4) &= f(x_4) + f(x_4 y_4) + f(y_4) = 0 + 2 + 2 = 4 \\ wt_f(x_5 y_5) &= f(x_5) + f(x_5 y_5) + f(y_5) = 0 + 3 + 2 = 5 \\ wt_f(x_6 y_6) &= f(x_6) + f(x_6 y_6) + f(y_6) = 2 + 2 + 2 = 6 \end{aligned}$$

It's form can be applied, so 3 is sufficient to become the lower bound of  $res(sun_3)$ . Then, we will prove for condition if  $n > 3$ . There are three cases for this condition.

Firstly if  $n \equiv 0 \pmod{3}$ . For this case we get,

$$\lceil \frac{n}{3} \rceil = \frac{n}{3}. \quad (2)$$

By (1), if  $n \equiv 0 \pmod{3}$  we get  $res(sun_n) \geq \lceil \frac{2n}{3} \rceil$ . Consequently by (2),

$$\begin{aligned} \lceil \frac{2n}{3} \rceil &= \frac{2n}{3} \\ &= 2 \lceil \frac{n}{3} \rceil. \end{aligned}$$

Secondly if  $n \equiv 1 \pmod{3}$ . For this case we get,

$$\lceil \frac{n}{3} \rceil = \frac{n-1}{3} + 1. \quad (3)$$

By (1), if  $n \equiv 1 \pmod{3}$  we get  $res(sun_n) \geq \lceil \frac{2n}{3} \rceil + 1$ . Consequently by (3),

$$\begin{aligned} \lceil \frac{2n}{3} \rceil + 1 &= \frac{2(n-1)}{3} + 1 + 1 \\ &= \frac{2(n-1)}{3} + 2 \\ &= 2\left(\frac{n-1}{3} + 1\right) \\ &= 2\lceil \frac{n}{3} \rceil. \end{aligned}$$

Thirdly if  $n \equiv 2 \pmod{3}$ ,

By (1), if  $n \equiv 2 \pmod{3}$  we get  $res(sun_n) \geq \lceil \frac{2n}{3} \rceil$ . For  $n \equiv 2 \pmod{3}$ ,  $\lceil \frac{2n}{3} \rceil$  has the same value as  $2\lceil \frac{n}{3} \rceil$ .

Next, we will prove the upper bound of  $res(sun_n)$ . To prove this part we construct  $k$ -labeling  $f$  with  $k = 3$  if  $n = 3$  and  $k = 2\lceil \frac{n}{3} \rceil$  if  $n > 3$  on  $sun_n$  as follows,

For  $n \geq 3$ ,

$$\begin{aligned} f(x_i) &= \begin{cases} 0, & \text{if } i = 1 \text{ and } 2, \\ \frac{4i-2}{3}, & \text{if } i = 3, 4, \dots, \lceil \frac{n}{2} \rceil \text{ and } i \equiv 2 \pmod{3}, \\ 4\lfloor \frac{i}{3} \rfloor, & \text{if } i = 3, 4, \dots, \lceil \frac{n}{2} \rceil \text{ and } i \not\equiv 2 \pmod{3}. \end{cases} \\ f(y_i) &= \begin{cases} \frac{4i-4}{3}, & \text{if } i = 1, 2, \dots, \lceil \frac{n}{2} \rceil \text{ and } i \equiv 1 \pmod{3}, \\ 2 + 4\lfloor \frac{i-2}{3} \rfloor, & \text{if } i = 1, 2, \dots, \lceil \frac{n}{2} \rceil \text{ and } i \not\equiv 1 \pmod{3}. \end{cases} \\ f(y_{n-(i-1)}) &= \begin{cases} \frac{4i}{3}, & \text{if } i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor \text{ and } i \equiv 0 \pmod{3}, \\ 2 + 4\lfloor \frac{i}{3} \rfloor, & \text{if } i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor \text{ and } i \not\equiv 0 \pmod{3}. \end{cases} \\ f(x_i y_i) &= \begin{cases} 2, & \text{if } i = 1 \text{ and } 2. \\ \frac{4i-6}{3}, & \text{if } i = 3, 4, \dots, \lceil \frac{n}{2} \rceil \text{ and } i \equiv 0 \pmod{3}, \\ 4\lfloor \frac{i}{3} \rfloor, & \text{if } i = 3, 4, \dots, \lceil \frac{n}{2} \rceil \text{ and } i \not\equiv 0 \pmod{3}. \end{cases} \\ f(x_1 x_n) &= 1. \end{aligned}$$

For  $n \geq 3$  and  $n \equiv 0 \pmod{6}$ ,

$$f(x_{n-(i-1)}) = \begin{cases} \frac{4i}{3}, & \text{if } i = \frac{n}{2}, \\ \frac{4i+4}{3}, & \text{if } i = 1, 2, \dots, (\frac{n}{2} - 1) \text{ and } i \equiv 2 \pmod{3}, \\ 2 + 4\lfloor \frac{i}{3} \rfloor, & \text{if } i = 1, 2, \dots, (\frac{n}{2} - 1) \text{ and } i \not\equiv 2 \pmod{3}. \end{cases}$$

For  $n \geq 3$  and  $n \not\equiv 0 \pmod{6}$ ,

$$f(x_{n-(i-1)}) = \begin{cases} \frac{4i+4}{3}, & \text{if } i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor \text{ and } i \equiv 2 \pmod{3}, \\ 2 + 4\lfloor \frac{i}{3} \rfloor, & \text{if } i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor \text{ and } i \not\equiv 2 \pmod{3}. \end{cases}$$

For  $n \geq 3$  and  $n \equiv 2 \pmod{6}$ ,

$$f(x_{n-(i-1)}y_{n-(i-1)}) = \begin{cases} \frac{4i-4}{3}, & \text{if } i = \frac{n}{2}, \\ \frac{4i+2}{3}, & \text{if } i = 1, 2, \dots, (\frac{n}{2} - 1) \text{ and } i \equiv 1 \pmod{3}, \\ 4\lceil \frac{i}{3} \rceil, & \text{if } i = 1, 2, \dots, (\frac{n}{2} - 1) \text{ and } i \not\equiv 1 \pmod{3}. \end{cases}$$

For  $n \geq 3$  and  $n \equiv 4 \pmod{6}$ ,

$$f(x_{n-(i-1)}y_{n-(i-1)}) = \begin{cases} \frac{4i-2}{3}, & \text{if } i = \frac{n}{2}, \\ \frac{4i+2}{3}, & \text{if } i = 1, 2, \dots, (\frac{n}{2} - 1) \text{ and } i \equiv 1 \pmod{3}, \\ 4\lceil \frac{i}{3} \rceil, & \text{if } i = 1, 2, \dots, (\frac{n}{2} - 1) \text{ and } i \not\equiv 1 \pmod{3}. \end{cases}$$

For  $n \geq 3$  and  $n \not\equiv 2, 4 \pmod{6}$ ,

$$f(x_{n-(i-1)}y_{n-(i-1)}) = \begin{cases} \frac{4i+2}{3}, & \text{if } i = 1, 2, \dots, (\lfloor \frac{n}{2} \rfloor) \text{ and } i \equiv 1 \pmod{3}, \\ 4\lceil \frac{i}{3} \rceil, & \text{if } i = 1, 2, \dots, (\lfloor \frac{n}{2} \rfloor) \text{ and } i \not\equiv 1 \pmod{3}. \end{cases}$$

For  $n = 3$ ,

$$f(x_i x_{i+1}) = 2i - 1, \text{ if } i = 1, 2.$$

For  $n > 3$  and  $n \equiv 3 \pmod{6}$ ,

$$f(x_i x_{i+1}) = \begin{cases} 1, & \text{if } i = 1 \text{ and } 2, \\ \frac{4i-5}{3}, & \text{if } i = \lceil \frac{n}{2} \rceil, \\ \frac{4i-9}{3}, & \text{if } i = 3, 4, \dots, (\lceil \frac{n}{2} \rceil - 1) \text{ and } i \equiv 0 \pmod{3}, \\ 3 + 4\lfloor \frac{i-4}{3} \rfloor, & \text{if } i = 3, 4, \dots, (\lceil \frac{n}{2} \rceil - 1) \text{ and } i \not\equiv 0 \pmod{3}. \end{cases}$$

For  $n > 3$  and  $n \not\equiv 3 \pmod{6}$ ,

$$f(x_i x_{i+1}) = \begin{cases} 1, & \text{if } i = 1 \text{ and } 2, \\ \frac{4i-9}{3}, & \text{if } i = 3, 4, \dots, (\lceil \frac{n}{2} \rceil) \text{ and } i \equiv 0 \pmod{3}, \\ 3 + 4\lfloor \frac{i-4}{3} \rfloor, & \text{if } i = 3, 4, \dots, (\lceil \frac{n}{2} \rceil) \text{ and } i \not\equiv 0 \pmod{3}. \end{cases}$$

For  $n \geq 3$  and  $n \equiv 0 \pmod{6}$

$$f(x_{n-(i-1)}x_{n-i}) = \begin{cases} \frac{4i+1}{3}, & \text{if } i = (\frac{n}{2} - 1), \\ \frac{4i-3}{3}, & \text{if } i = 1, 2, \dots, (\frac{n}{2} - 2) \text{ and } i \equiv 0 \pmod{3}, \\ 1 + 4\lfloor \frac{i}{3} \rfloor, & \text{if } i = 1, 2, \dots, (\frac{n}{2} - 2) \text{ and } i \not\equiv 0 \pmod{3}. \end{cases}$$

For  $n \geq 3$  and  $n \not\equiv 0 \pmod{6}$

$$f(x_{n-(i-1)}x_{n-i}) = \begin{cases} \frac{4i-3}{3}, & \text{if } i = 1, 2, \dots, (\lfloor \frac{n}{2} \rfloor - 1) \text{ and } i \equiv 0 \pmod{3}, \\ 1 + 4\lfloor \frac{i}{3} \rfloor, & \text{if } i = 1, 2, \dots, (\lfloor \frac{n}{2} \rfloor - 1) \text{ and } i \not\equiv 0 \pmod{3}. \end{cases}$$

Then, by  $f$ -labeling we know that label of vertices is even positive integer. So the upper bound of  $res(sun_n)$  has same value as the lower bound of  $res(sun_n)$ . The weight of edges are, For even  $n$ ,

$$\begin{aligned}
 wt(x_i y_i) &= \begin{cases} 2, & \text{if } i = 1, \\ 4(i - 1), & \text{if } i = 2, 3, \dots, (\frac{n}{2} + 1). \end{cases} \\
 wt(x_{n-(i-1)} y_{n-(i-1)}) &= 4i + 2, \text{ if } i = 1, 2, \dots, (\frac{n}{2} - 1). \\
 wt(x_i x_{i+1}) &= 4i - 3, \text{ if } i = 1, 2, \dots, \lceil \frac{n}{2} \rceil. \\
 wt(x_{n-(i-1)} x_{n-i}) &= 4i + 3, \text{ if } i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor. \\
 wt(x_1 x_n) &= 3, \text{ if } n \geq 3.
 \end{aligned}$$

For odd  $n$ ,

$$\begin{aligned}
 wt(x_i y_i) &= \begin{cases} 2, & \text{if } i = 1, \\ 4(i - 1), & \text{if } i = 2, 3, \dots, \lceil \frac{n}{2} \rceil. \end{cases} \\
 wt(x_{n-(i-1)} y_{n-(i-1)}) &= 4i + 2, \text{ if } i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor. \\
 wt(x_i x_{i+1}) &= 4i - 3, \text{ if } i = 1, 2, \dots, \lceil \frac{n}{2} \rceil. \\
 wt(x_{n-(i-1)} x_{n-i}) &= 4i + 3, \text{ if } i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor. \\
 wt(x_1 x_n) &= 3, \text{ if } n \geq 3.
 \end{aligned}$$

By this investigate, we can conclude that every edges has different weight, consequently  $f$  is edge irregular reflexive  $k$ -labeling. So, the proof of  $res(sun_n)$  is completed.  $\square$

An illustration of edge irregular reflexive  $k$ -labeling on  $sun_n$  for even  $n$  can be seen on Figure 1. The black color is label, red color is weight and blue color is name of vertices.




Figure 1. Edge irregular reflexive 4-labeling of  $sun_4$ .

### 3. The corona of cycle and null graph with two vertices

According Dwivedi [4], a graph  $G(V, E)$  is called null graph which is does not have any edges, in other words every vertex are isolated, denoted as  $N_m$  with  $m$  the number of vertices. Then, based on definition corona of two graph by Harry and Frunct [3], corona of cycle and null graph denoted as  $C_n \odot N_m$  is a graph that formed by one copy of graph  $C_n$  and  $n$ -copy graph of  $N_m$  with  $i$  - th vertex from  $C_n$  is connected to all of vertices from  $i$  - th copy of graph  $N_m$ . It's consequently  $C_n \odot N_m$  have  $|E| = n(m + 1)$ . First, we know that  $C_n \odot N_1$  same as sun graph. Therefore, in this paper we continue to discuss of  $C_n \odot N_2$  for  $C_n$  with  $n \geq 3$ . We will denote vertices in the cycle of  $C_n \odot N_2$  as  $x_i$  and pendant vertices denoted as  $y_{i,j}$ . So set of vertices  $V(C_n \odot N_2) = \{x_i, y_{i,j} : 1 \leq i \leq n \text{ and } j = 1, 2\}$ , consequently set of edges  $E(C_n \odot N_2) = \{x_i x_{i+1}, x_i y_{i,j} : 1 \leq i \leq n \text{ and } j = 1, 2\}$ . Then reflexive edge strength of  $C_n \odot N_2$  can be found on *Theorem 3.1*.

**Theorem 3.1.** For  $C_n \odot N_2$  with  $n \geq 3$ ,

$$res(C_n \odot N_2) = 2 \lfloor \frac{n+1}{2} \rfloor, \text{ if } n \geq 3.$$

*Proof.* First, we prove the lower bound of  $res(C_n \odot N_2)$ . By section (3) we get  $|E|$  of  $C_n \odot N_2$  is  $3n$ , then by *Lemma 1.1* we get

$$res(C_n \odot N_2) \geq \begin{cases} \lceil \frac{3n}{3} \rceil, & \text{if } 3n \not\equiv 2, 3 \pmod{6}, \\ \lceil \frac{3n}{3} \rceil + 1, & \text{if } 3n \equiv 2, 3 \pmod{6}. \end{cases} \tag{4}$$

It's equivalent with,

$$res(C_n \odot N_2) \geq \begin{cases} n, & \text{if } 3n \not\equiv 2, 3 \pmod{6}, \\ n + 1, & \text{if } 3n \equiv 2, 3 \pmod{6}. \end{cases} \tag{5}$$

For  $n \equiv 0 \pmod{2}$  we get,

$$\lfloor \frac{n+1}{2} \rfloor = \frac{n}{2}. \tag{6}$$

By (5), if  $n \equiv 0 \pmod{2}$  we get  $res(C_n \odot N_2) \geq n$ . Consequently by (6) we get,

$$\begin{aligned} n &= \frac{2n}{2} \\ &= 2 \lfloor \frac{n+1}{2} \rfloor. \end{aligned}$$

For  $n \equiv 1 \pmod{2}$  we get,

$$\lfloor \frac{n+1}{2} \rfloor = \frac{n+1}{2}. \tag{7}$$

By (5), if  $n \equiv 1 \pmod{2}$  we get  $res(C_n \odot N_2) \geq n + 1$ . Consequently by (7) we get,

$$\begin{aligned} n + 1 &= 2 \left( \frac{n+1}{2} \right) \\ &= 2 \lfloor \frac{n+1}{2} \rfloor. \end{aligned}$$

Next, we will prove the upper bound of  $res(C_n \odot N_2)$  for  $n \geq 3$ . To prove this part we construct  $k$ -labeling  $f$  with  $k = 2\lfloor \frac{n+1}{2} \rfloor$  if  $n \geq 3$  on  $C_n \odot N_2$  as follows,

For  $n \geq 3$ ,

$$\begin{aligned} f(x_i) &= \begin{cases} 2(i-1), & \text{if } i = 1 \text{ and } 2, \\ 2i, & \text{if } i = 3, 4, \dots, \lceil \frac{n}{2} \rceil. \end{cases} \\ f(x_{n-(i-1)}) &= 2i, \text{ if } i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor. \\ f(y_{i,j}) &= \begin{cases} 0, & \text{if } i = 1 \text{ and } j = 1, 2, \\ 2i, & \text{if } i = 2, 3, \dots, \lceil \frac{n}{2} \rceil \text{ and } j = 1, 2. \end{cases} \\ f(y_{(i-1),j}) &= 2i, \text{ if } i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor \text{ and } j = 1, 2. \\ f(x_{n-(i-1)}x_{n-i}) &= 2i + 1, \text{ if } i = 1, 2, \dots, (\lfloor \frac{n}{2} \rfloor - 1). \\ f(x_{n-(i-1)}y_{n-(i-1)}) &= 2i + j - 2, \text{ if } i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor \text{ and } j = 1, 2. \\ f(x_1x_n) &= 1. \end{aligned}$$

For  $n = 3$ ,

$$f(x_iy_{i,j}) = \begin{cases} j, & \text{if } i = 1, \\ j + 1, & \text{if } i = 2. \end{cases}$$

For  $n > 3$  and  $n \equiv 0 \pmod{2}$ ,

$$f(x_iy_{i,j}) = \begin{cases} j, & \text{if } i = 1, 2 \text{ and } j = 1, 2, \\ 2i + j - 6, & \text{if } i = 3, 4, \dots, \lceil \frac{n}{2} \rceil \text{ and } j = 1, 2. \end{cases}$$

For  $n > 3$  and  $n \equiv 1 \pmod{2}$ ,

$$f(x_iy_{i,j}) = \begin{cases} j, & \text{if } i = 1, 2 \text{ and } j = 1, 2, \\ 2i + j - 6, & \text{if } i = 3, 4, \dots, \lfloor \frac{n}{2} \rfloor \text{ and } j = 1, 2, \\ 2i + j - 5, & \text{if } i = \lceil \frac{n}{2} \rceil \text{ and } j = 1, 2. \end{cases}$$

For  $n \geq 3$  and  $n \equiv 0 \pmod{2}$ ,

$$f(x_ix_{i+1}) = \begin{cases} 2, & \text{if } i = 1, \\ 4, & \text{if } i = 2 \text{ and } n = 4, \\ 2, & \text{if } i = 2 \text{ and } n \neq 4, \\ 2i - 4, & \text{if } i = 3, 4, \dots, (\frac{n}{2} - 1) \text{ and } n > 4, \\ 2i - 2, & \text{if } i = \frac{n}{2} \text{ and } n > 4. \end{cases}$$



For  $n \geq 3$  and  $n \equiv 1 \pmod{2}$ ,

$$f(x_i x_{i+1}) = \begin{cases} 2, & \text{if } i = 1, \\ 3, & \text{if } i = 2 \text{ and } n = 3, \\ 2, & \text{if } i = 2 \text{ and } n \neq 3, \\ 2i - 4, & \text{if } i = 3, 4, \dots, (\lceil \frac{n}{2} \rceil - 1) \text{ and } n > 3, \\ 2i - 3, & \text{if } i = \lceil \frac{n}{2} \rceil \text{ and } n > 3. \end{cases}$$

Then, by  $f$ -labeling we know that label of vertices is even positive integer. From the above formula, we get that the upper bound of  $res(C_n \odot N_2)$  same as the lower bound and the weight of edges we get,

For even  $n$ ,

$$\begin{aligned} wt(x_i y_{i,j}) &= 6i + j - 6, \text{ if } i = 1, 2, \dots, \frac{n}{2} \text{ and } j = 1, 2. \\ wt(x_{n-(i-1)} y_{n-(i-1),j}) &= 6i + j - 2, \text{ if } i = 1, 2, \dots, \frac{n}{2} \text{ and } j = 1, 2. \\ wt(x_i x_{i+1}) &= 6i - 2, \text{ if } i = 1, 2, \dots, \frac{n}{2}. \\ wt(x_{n-(i-1)} x_{n-i}) &= 6i + 3, \text{ if } i = 1, 2, \dots, (\frac{n}{2} - 1). \\ wt(x_1 x_n) &= 3, \text{ if } n \geq 3. \end{aligned}$$

For odd  $n$ ,

$$\begin{aligned} wt(x_i y_{i,j}) &= \begin{cases} 6i + j - 6, & \text{if } i = 1, 2, \dots, (\lceil \frac{n}{2} \rceil - 1) \text{ and } j = 1, 2, \\ 6i + j - 5, & \text{if } i = \lceil \frac{n}{2} \rceil \text{ and } j = 1, 2. \end{cases} \\ wt(x_{n-(i-1)} y_{n-(i-1),j}) &= 6i + j - 2, \text{ if } i = 1, 2, \dots, \lfloor \frac{n}{2} \rfloor \text{ and } j = 1, 2. \\ wt(x_i x_{i+1}) &= \begin{cases} 6i - 2, & \text{if } i = 1, 2, \dots, (\lceil \frac{n}{2} \rceil - 1), \\ 6i - 5, & \text{if } i = \lceil \frac{n}{2} \rceil. \end{cases} \\ wt(x_{n-(i-1)} x_{n-i}) &= 6i + 3, \text{ if } i = 1, 2, \dots, (\lfloor \frac{n}{2} \rfloor - 1). \\ wt(x_1 x_n) &= 3, \text{ if } n \geq 3. \end{aligned}$$

By this investigate, we can conclude that every edges has different weight, consequently  $f$  is edge irregular reflexive  $k$ -labeling. So, the proof of  $res(C_n \odot N_2)$  is completed.  $\square$

An illustration of edge irregular reflexive  $k$ -labeling on  $C_n \odot N_2$  can be seen on Figure 2. The black color is label, red color is weight and blue color is name of vertices.




Figure 2. Edge irregular reflexive 4-labeling of  $C_4 \odot N_2$ .

#### 4. Concluding remark

As result the discussion we get conclude  $res(sun_n)$  or  $res(C_n \odot N_1)$  are 3 for  $n = 3$  and  $2\lceil \frac{n}{2} \rceil$  for  $n > 3$ , while  $res(C_n \odot N_2)$  is  $2\lfloor \frac{n+1}{2} \rfloor$  for  $n \geq 3$ . Moreover, there is open problem for next research about this graph, which still on progress to investigate.

**Open problem:** What is reflexive edge strength of  $C_n \odot N_m$  for  $n \geq 3$  and  $m \geq 3$ .

#### Acknowledgement

This research in this article was supported by RKAT PTNBH Universitas Sebelas Maret fiscal year 2021 through scheme research of grant research group (hgr-uns research) with contract number 260/UN27.22/HK.07.00/2021.

#### References

- [1] D. Indriati, Widodo, and I. Rosyida, Edge irregular reflexive labeling on corona of path and other graphs. *J.P. Conf. Ser.*, **1489** (2020), 012004.
- [2] D. Tanna, J. Ryan, and A. Semaničová-Feňovčíková, Edge irregular reflexive labeling of prisms and wheels, *Australas. J. Combin.*, **69(3)** (2017), 394–901.
- [3] F. Harary and R. Frucht, On the corona of two graphs, *Aequationes Math.*, **4** (1970), 322–325.
- [4] G.K. Dwivedi, *Analysis and Desain of Algorithm*, Laxmi Publications (P) LTD, New Delhi (2007).
- [5] J.A. Galian, A Dynamic survey of graph labeling, *Electron. J. Combin.*, **20** (2017), #DS6.
- [6] J. Ryan, B. Munasinghe, D. Tanna, Reflexive irregular labelings, *Unpublished work*, 2017.
- [7] M. Bača, M. Irfan, J. Ryan, A. Semaničová-Feňovčíková, and D. Tanna, Note on edge irregular reflexive labelings of graphs, *AKCE Int. J. Graphs Comb.*, **16** (2010), 145–157.
- [8] M. Bača, S. Jendrol', M. Miller, and J. Ryan, On irregular total labeling, *Discrete Math.*, **307** (2007), 1378–1388.

- [9] R. Boulet, Spectral characterizations of sun graphs and broken sun graphs, *Discrete Math. Theor. Comput. Sci.*, **11:2** (2009), 149–160.
- [10] W.D. Wallis and A.M. Marr, *Magic Graphs Second edition*, Springer Science and Business Media, New York (2013).